

# High $p_T$ hadron suppression in Au+Au collisions at RHIC due to final-state interactions

B.J. Choi<sup>1</sup>, D.H. Hardtke<sup>1</sup>, P.M. Jacobs<sup>1</sup>, J.L. Klay<sup>1</sup>

<sup>1</sup>Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720




FIG. 1: Nuclear modification factor  $R_{AB}(p_T)$  in central Au+Au collisions (blue stars) and minimum bias and central d+Au collisions at  $\sqrt{s_{NN}}=200\text{GeV}$ .

Charged hadron production at high  $p_T$  in central Au+Au collisions at RHIC is highly suppressed compared to the expectation for a superposition of incoherent inelastic N+N collisions[1, 2]. We have also observed that away-side jet correlations are suppressed in central Au+Au collisions while the near-angle correlations in all centralities of Au+Au are strikingly similar to those in p+p collisions at the same energy[3]. These effects could be attributed to gluon saturation in the initial nuclear wavefunction which suppresses the production rate for di-jets (“initial-state”)[4] or to energy loss of the partner jet as it bores through the medium created in a heavy ion collision (“final-state” effect)[5, 6].

If the observed “jet-quenching” is due to initial-state nuclear effects, one would expect to see a similar suppression of hadron production and di-jet correlations in d+Au collisions at  $\sqrt{s_{NN}}=200\text{GeV}$ . If the observations are primarily due to final-state effects, one would expect comparable results in p+p and d+Au collisions, where no extended nuclear medium is produced. During 2003, RHIC undertook an extended run with d+Au collisions. We analyzed the data from these collisions using the STAR detector and compared the results on high  $p_T$  hadron production and di-jet correlations in p+p, d+Au and Au+Au collisions at  $\sqrt{s_{NN}}=200\text{GeV}$ .

Figure 1 shows the “nuclear modification factor”,  $R_{AB}(p_T)$ , which is the ratio of charged hadron production in central Au+Au collisions and two centralities of d+Au collisions compared to the  $\langle N_{\text{bin}} \rangle$ -scaled spectrum in p+p collisions. In d+Au collisions, there is an approximately 20% enhancement of charged hadron production at intermediate  $p_T$  consistent with expectations from the Cronin effect, in contrast to the factor of five suppression in central Au+Au collisions.

Figure 2 shows the azimuthal correlations between a high  $p_T$  trigger particle with  $4 < p_T < 6\text{ GeV/c}$  and associated parti-




FIG. 2: Di-hadron correlations in central Au+Au collisions (blue stars), central d+Au collisions (red circles) and p+p collisions (black histogram) at  $\sqrt{s_{NN}}=200\text{GeV}$ .

cles with  $2 < p_T < 4\text{ GeV/c}$  in p+p, d+Au and Au+Au collisions at  $\sqrt{s_{NN}}=200\text{GeV}$ . In all three cases the near-side correlations have similar shape and magnitude, consistent with standard jet fragmentation. The away-side correlations in p+p and d+Au collisions are very similar, while for central Au+Au collisions the away-side correlations are completely suppressed.

These results demonstrate conclusively that the striking suppression phenomena observed in central Au+Au collisions are not due to gluon saturation in the initial-state nuclear wavefunction, but are rather due to the interaction of high energy partons or their fragmentation products in the dense medium created in such collisions[7].

- 
- [1] C. Adler et al. (STAR Collaboration), Phys. Rev. Lett. **89**, 202301 (2002).
  - [2] J. Adams et al. (STAR Collaboration), Phys. Rev. Lett. **91**, 172302 (2003).
  - [3] C. Adler et al., Phys. Rev. Lett. **90**, 082302 (2003).
  - [4] D. Kharzeev, E. Levin, and L. McLerran, Phys. Lett. B **561**, 93 (2003).
  - [5] R. Baier, D. Schiff, and B. G. Zakharov, Ann. Rev. Nucl. Part. Sci. **50**, 37 (2000).
  - [6] M. Gyulassy, I. Vitev, X. N. Wang, and B. Zhang (2003), nucl-th/0302077.
  - [7] J. Adams et al. (STAR Collaboration), Phys. Rev. Lett. **91**, 072304 (2003).